



Biotoxin Illness Survey

PLEASE INDICATE WHETHER YOU ARE EXPERIENCING ANY OF THE FOLLOWING SYMPTOMS

- Weak
- Decreased Assimilation of New Knowledge
- Aches
- Headache
- Light Sensitivity

Fatigue

- Memory Impairment
- Decreased Word Finding

Difficulty Concentrating

- Joint Pain
- A.M. Stiffness
- Cramps

- Unusual Skin Sensitivity
- Tingling

- Shortness of Breath
- Sinus Congestion

- Appetite Swings
- Difficulty Regulating Body Temperature
- Increased Urinary Frequency

- Red Eyes
- Blurred Vision
- Sweats (night)
- Mood Swings
- Ice-pick Pain

- Abdominal Pain
- Diarrhea
- Numbness

- Tearing of Eyes
- Disorientation
- Metallic Taste

- Static Shocks
- Vertigo

- Cough
- Excessive Thirst
- Confusion

Biotoxin Illness Survey Assessment

The Biotoxin Illness Survey uses 35 specific symptoms, divided into 13 groups/clusters.

Examine your client's survey responses and count how many groups have at least one checked box. According to Dr. Shoemaker's criteria, if a client has symptoms in eight or more clusters, they have a high likelihood of biotoxin illness.

If their visual contrast sensitivity (VCS) test results are also positive, the chances that they have biotoxin illness are greater than 99 percent.

**Based on the biotoxin symptom clusters developed by Dr. Ritchie Shoemaker of SurvivingMold.com.*